

0759
3795

BULLETIN

39 1/2

DE LA

SOCIÉTÉ IMPÉRIALE

DES NATURALISTES

DE MOSCOU.

TOME LIV.

ANNÉE 1879.



№ 1.

MOSCOU.

Imprimerie de l'Université Impériale.
(M. Katkoff.)

1879.

N 4967

ribus, solitoque paulo minus pilosis (lamina 15 : 30 vel 20 : 40 mill. magna), fructus apice tantum pilosi.

Prunus pogonostyla Maxim. (*Cerasus*) fruticosa humilis, ramulis floriferis dense pubescentibus, foliis hysteranthis brevissime petiolatis ovatis subito acuminatis, superne adpresse parcius, subtus ad reticulum patentim rufopilosis duplicato-serratis venis immersis, stipulis petiolo longioribus setaceis glandulosociliatis basi laciniatis, pedunculis 1—2 dense pilosis brevissimis, fructiferis $\frac{1}{2}$ —1-pollicaribus; calycis utrinque pubescentis tubo late campanulato brevioris quam lacinae ovato-oblongae glanduloso-serratae, petalis obovatis calycem duplo superantibus, stylo stamina aequante infra medium longe paleaceopiloso, ovario glabro, drupa (nondum matura) styli basi hirsuta apiculata ovaliglobosa, putamine laevi hinc 1- illic 2-sulcato. *Prunus spec.* Hance in Trim. Journ. of bot. 1875, quoad pl. Grijsi).

In summo cacumine montis Nam-tau-wú, prope Amoy, alt. circa 2000 ped., Decembri 1862 fl. (de Grijs in hb. Hance n. 10,130.), prope Tamsuy ins. Formosae (Oldham n. 105. frf., 1864).

A proxima *P. humili* Bge differt foliorum forma et pube, et praesertim stylo basi hispido. Folia fere *Armeniaca sibirica* forma, sed brevius cuspidata. Pedunculi deflorati sensim elongantur.

Prunus humilis Bge Enum. Chin. n. 133.

In China boreali vulgaris: circa Pekinum (Bunge, Kirilow), Lung-tsüan-sze fine Maji fr. juv., Miao-feng-shan, fine Augusti fr. mat., Takio-sze, optime evoluta altit. 3—4 mill. pedum, fine Aug. fructif., Po-hua-shan, flor. (Bretschneider). In Mongolia maxime australi, circa Kalgan (Ladyshinski).