

Repertorium specierum novarum regni vegetabilis.

**Centralblatt für Sammlung und Veröffentlichung
von Einzeldiagnosen neuer Pflanzen.**

Herausgegeben

von

Dr. phil. Friedrich Fedde,
Herausgeber von Just's Botanischem Jahresberichte

Fasciculus VIII (1910).

BERLIN-WILMERSDORF.

**SELBSTVERLAG DES HERAUSGEBERS, WEIMARISCHE STR. 3^E
KOMMISSIONS-VERLAG VON GEBRÜDER BORNTRAEGER, BERLIN.**

— 1910 —

Repertorium specierum novarum regni vegetabilis

auctore

F. Fedde

No. 182/184

VIII. Band

10. August 1910

CXIII. Decades plantarum novarum. XXXIX.

Auctore H. Léveillé.

(Originaldiagnosen.)

381. **Blumea velutina** Lévl. et Vant. nomen novum. (*Senecio velutinus* Lévl. et Vant. in Fedde, Repertorium, VI (1909), p. 331.

382. **Blumea compactiflora** Lévl. et Vant., nov. spec.

Affinis *B. hieracifolia*; a quo tamen differt rosula radicali duplice, heterophylla et ramulis novellis diversifoliis; foliis spatulatis, alato-petiolatis et crenatis folia novella lanceolata vix vel non dentata sunt intermixta; capitula capitata compacta.

Kouy-Tchéou: Environs de Ma-Jo sur la droite, mars 1909, 20 avril 1908; no. 3690; Lo-Fou cultures, avril 1909 (Jul. Cavalerie.)

383. **Saussurea Cavaleriei** Lévl. et Vant., nov. spec.

Planta robusta, puberula, rubescens; caulis striatus; folia petiolata, late ovata, argute dentata, caudato-acuminata (inferiora non visa); inflorescentia diffusa, bracteata, bracteis lanceolato-acuminatis; capitula cylindrica bina; involucra propter bracteas paleaceas apice nigras et reflexas variegata; flores medii pulchre atro-purpureo-violacei; tubo flosculi limbo alte fisso et lobis linearibus paulo longiore; tubo staminali longe exerto; akoenia nigra, tetragona, conico-cylindrica, glabra, pappo ferrugineo.

Kouy-Tchéou: Oct. 1908 (Jul. Cavalerie, 2976).

384. **Scutellaria glechomaefolia** Lévl. et Vant., nov. spec.

Planta pubescens; caulis simplex; folia remota, longe petiolata, corodata illa *Glechomae hederaceae* perfecte referentia; flores pro genere mediocres, terminales, hirtelli; calyx hirsutus; corolla calice 8-plo longior.

Corea: Quelpaert, in herbidis, jun. 1907 (Faurie, 1929).

385. **Scutellaria multibrachiata** Lévl. et Vant., nov. spec.

Affinis praecedenti cuius forsitan varietas; a quo tamen differt sequentibus notis: caule a basi ramosissimo; statura, foliis et floribus minoribus.

Corea: Quelpaert, in silvis Hallaisan, juin. 1909 (Taquet, 3088).

386. **Scutellaria Fauriei** Lévl. et Vant., nov. spec.

Planta pubescens, grisea, humilis et repens; folia parvula, obtusa, crenata, petiolata, inferne rotundata; flores pro statura plantae magni;

inflorescentia spicata, elongata, terminalis, bracteolata; corolla calice 10-plo longior.

Corea: Quelpaert, in silvis, jun. 1907 (Faurie, 1943).

387. Scutellaria Komarovii Lévl. et Vant., nov. spec.

Planta valde elata erecta, macilenta, puberula, a basi longe sed pauciramosa; folia distantia, rara, angusta, sessilia, dentata; flores pro genere mediocres, puberuli, bracteati; corolla calice 7—8 plo longior.

Corea: In humidis Taikou, 30 maio 1906 (Faurie, 819).

388. Scutellaria Taquetii Lévl. et Vant., nov. spec.

Planta longissime repens; radice articulata sed non inflata, caule viloso, infra nudo, superne dense folioso; folia villosa illa *Teucrui cha-maedrys* referentia, sed sessilia; inflorescentia axillaris; flores pro genere mediocres; corolla calice 5—6 longior.

Corea: Quelpaert, in herbidis Tpyengeni, juin. 1909 (Taquet, 3089).

389. Scutellaria Esquirolii Lévl. et Vant., nov. spec.

Planta speciosa, dumosa ramosissima, puberula; inflorescentia ramosa terminalis; folia ovata, acuta, integra; flores et fructus rosei; flores pro genere mediocres; corolla calice 4-plo longior.

Kouy-Tchéou: Depuis le col de Pay-So jusqu'à Pai-Yang par Kay-Lang, 22 juin 1905 (Jos. Esquirol, 484).

390. Scutellaria Cavaleriei Lévl. et Vant., nov. spec.

Planta ramosa, caulis glabris; folia sessilia ovata, acuta, crenata subtus in nervis hirtella, supra granulata; inflorescentia terminalis, puberula, elongata, bracteolata; flores violacei, pro genere mediocres; corolla calice 6—8-plo longior.

Kouy-Tchéou: Gan-Pin, 10 mai 1904 (Jul. Cavalerie, 2105); Kiang-Sou (d'Argy).

CXIV. Alfred J. Ewart, *Contributiones Florae Australiensis. I.*

(Ex: Proc. Roy. Soc. Victoria N. S. XIX, 2 [1907], pp. 33—45, pl. X—XII.)

Bellida nov. gen. (*Compositae*) A. J. Ewart, l. c., p. 34.

Annuals or perennials with radical leaves, and inflorescences at the ends of long simple stalks, leafless, or with a small bract on the shaft. Bracts of the involucre imbricate, scarious in a double series, the inner ones larger. Florets all alike, regularly 5-toothed, tubular and yellow, with no scales between them. Anthers with well-marked appendices and with rounded bases. Style and stigma of Asterae. Fruit on a distinct stalk, and obliquely inserted on the head. Pappus of two small, separable, cup-like scales, each bearing a single row of stiff bristles.

Relationships. — The obliquely lateral insertion of the fruit reminds one of the *Cynareae-Centaurineae*, from which the plant differs widely in other respects. The character of the bracts and the homogamous inflorescences suggest the *Gnaphaleae*, but the anthers have rounded bases and no tails.